


STANDING ORDERS FOR CHEST X-RAYS

DATE: March 11, 2016

TO: WV-DTBE Radiological Technicians

FROM: Matthew Simmons, MD  2-11-16
WV Division of TB Elimination

1. Evaluate with a posterior-anterior (PA) chest radiograph anyone presenting to Chest X-Ray or Chest Diagnostic Clinic who has the following:
 - Newly recognized positive protein derivative (PPD) result
 - Newly recognized positive Interferon Gamma Release Assay (IGRA) result
 - PPD test conversion
 - IGRA conversion
 - Symptoms suggestive of active TB (cough lasting longer than two weeks, chest pain, hemoptysis, fatigue, weight loss, poor/no appetite, chills, fever, night sweats)
 - Being monitored regularly by serial chest x-rays in Chest Diagnostic Clinic
2. Provide proper radiation protection to patient as well as technologist as per radiologic standards (i.e., shielding of child-bearing age, gonadal protection, etc.)
3. Do not do radiographs on pregnant women. Consult with physician. If x-ray is done with physician order, ensure shielding is shown on x-ray film and proper documentation is in patient chart
4. Use appropriate markers on all radiographs

**Bureau for Public Health
Office of Epidemiology and Prevention Services
Division of Tuberculosis Elimination**
350 Capitol Street, Room 125
Charleston, West Virginia 25301
(304) 558-3669